arXiv:1602.06069v1 [math.NT] 19 Feb 2016

A NOTE ON THE GAPS BETWEEN ZEROS OF EPSTEIN'S ZETA-FUNCTIONS ON THE CRITICAL LINE

STEPHAN BAIER, SRINIVAS KOTYADA AND USHA KESHAV SANGALE

Abstract. It is proved that Epstein's zeta-function $\zeta_Q(s)$, related to a positive definite integral binary quadratic form, has a zero $1/2+i\gamma$ with $T \le \gamma \le T+T^{3/7+\varepsilon}$ for sufficiently large positive numbers T. This is an improvement of the result by M. Jutila and K. Srinivas (Bull. London Math. Soc. 37 (2005) 45–53).

To Professor Matti Jutila with deep regards

1. Introduction

Let the quadratic form $Q(x,y) = ax^2 + bxy + cy^2$ be positive definite, has integer coefficients and let $r_Q(n)$ count the number of solutions of the equation $Q(x,y) = n$ in integers x and y. The Epstein zeta-function associated to Q is denoted by $\zeta_Q(s)$ and is given by the series

$$\zeta_Q(s) = \sum_{(x,y)\in\mathbb{Z}^2-(0,0)} \frac{1}{Q(x,y)^s} = \sum_{n=1}^{\infty} \frac{r_Q(n)}{n^s}$$

in the half-plane $\sigma > 1$, where (as usual) $s = \sigma + it$. Throughout the paper, we shall write Δ to denote the number $\Delta = 4ac - b^2 > 0$. $\zeta_Q(s)$ has many analytical properties in common with the Riemann zeta-function, $\zeta(s)$. For example, it admits analytic continuation into the entire complex plane except for a simple pole at $s = 1$ with residue $2\pi\Delta^{-1/2}$. It satisfies the following functional equation

$$\left(\frac{\sqrt{\Delta}}{2\pi}\right)^s \Gamma(s)\zeta_Q(s) = \left(\frac{\sqrt{\Delta}}{2\pi}\right)^{1-s} \Gamma(1-s)\zeta_Q(1-s) \tag{1.1}$$

The analogue of Hardy's theorem for $\zeta(s)$ also holds true for $\zeta_Q(s)$, i.e., $\zeta_Q(s)$ admits infinitely many zeros on the critical line $\sigma = 1/2$. In fact, much more is true. In 1934, Potter and Tichmarsh [3] showed that every interval of the type $[T, T+T^{1/2+\varepsilon}]$ contains a zero $1/2+i\gamma$ of $\zeta_Q(s)$ for any fixed ε and for all sufficiently large T. Sankaranarayan in 1995 [4] showed that the same result holds true for intervals of the type $[T, T+cT^{1/2}\log T]$. In 2005, Jutila and Srinivas [2] proved that the same is true for intervals of the type $[T, T+cT^{5/11+\varepsilon}]$, thus surpassing the classical barrier of $1/2$ in exponent of T. In this note we improve this result further. More precisely, we prove the following

Theorem 1. *Let Q be a positive definite binary integral quadratic form. Then for any fixed $\varepsilon > 0$ and $T \ge T(\varepsilon, Q)$, there is a zero $1/2+i\gamma$ of the corresponding Epstein zeta-function $\zeta_Q(s)$ with*

$$|\,\gamma - T\,| \le T^{3/7+\varepsilon}. \tag{1.2}$$

Our exposition closely follows the presentation and technical details given in [2], therefore, in this note we indicate only those steps in [2] which enable us to improve the result of Jutila and Srinivas mentioned earlier.

The rest of the paper is organized as follows: in section 2 we briefly discuss the main exponential sum of [2], section 3 contains basic results used in the proof of the main theorem and in section 4 we estimate the double exponential sum non-trivially, which leads to the improvement.

2000 *Mathematics Subject Classification.* 11E45 (primary); 11M41 (secondary).

2. The critical exponential sum

In order to show that $\zeta_Q(s)$ has a zero $1/2+i\gamma$ with $|T-\gamma|\le H$, the main point in [2] was to show that the sum

$$\sum_n \eta(n) r_Q(n) n^{-1/2-it} \tag{2.1}$$

is *small* in a certain sense, where $\eta(n)$ is a smooth weight function supported in the interval $[T\sqrt{\Delta}/2\pi - K, T\sqrt{\Delta}/2\pi + K]$, t lies *close* to T, H and K are parameters related by $HK = T^{1+\varepsilon}$. To estimate the sum (2.1), it was first transformed into another sum (equation (3.6) of [2]) using a transformation formula. By partial summation, this sum got reduced to the estimation of the following term:

$$K^{1/4}N^{-1/4}T^{-1/2} \sum_{N\le Q^*(x,y)\le N'} e\left(Q^*(x,y)\left(\frac{\overline{h\Delta_0}}{k} - \frac{1}{2hk\Delta_0}\right) + \frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right)\right) \tag{2.2}$$

This is equation (4.4) of [2] (see 4 for details). The objective was to show that (2.2) is $O(T^{-\varepsilon})$. To achieve this, the double exponential sum, appearing above, was estimated non-trivially in one variable using van der Corput's method and trivial estimate was taken in the other variable. They obtained the bound $O(K^{11/12}T^{-1/2})$, which is $O(T^{-\varepsilon})$, provided $K = T^{6/11-\varepsilon}$.

In the present paper we estimate the above double exponential sum non-trivially in both variables. To show that (2.2) is $O(T^{-\varepsilon})$, it is now enough to take $K = T^{4/7-\varepsilon}$ and thus the gap $H = T^{3/7+\varepsilon}$ follows.

3. Preliminary lemmas

We will use the following well-known lemmas in the proof of our theorem.

Lemma 1. *(Generalized Weyl differencing) Let $a<b$ be integers, λ be a natural number and $\xi(n)$ be a complex valued function such that $\xi(n)=0$ if $n\notin(a,b]$. If H is a positive integer then*

$$|\sum_n \xi(n)|^2 \le \frac{(b-a)+H}{H}\sum_{|h|<H}\left(1-\frac{\lambda|h|}{H}\right)\sum_n \xi(n)\overline{\xi(n-\lambda h)} \tag{3.1}$$

Proof. This is Lemma 2.5 of [1] for the case $\lambda = 1$. The general case follows similarly.

Lemma 2. *(B-Process , Lemma 3.6 of [1]) Suppose that f has four continuous derivatives on $[a,b]$, and that $f''<0$ on this interval. Suppose further that $[a,b]\subseteq[N,2N]$ and that $\alpha = f'(b)$ and $\beta = f'(a)$. Assume that there is some $F>0$ such that*

$$f^{(2)}(x) \asymp FN^{-2},\ f^{(3)}(x) \ll FN^{-3}, \text{and } f^{(4)}(x) \ll FN^{-4}$$

for x in $[a,b]$. Let x_ν be defined by the relation $f'(x_\nu)=\nu$, and let $\phi(\nu) = -f(x_\nu)+\nu x_\nu$. Then

$$\sum_{a\le n\le b} e(f(n)) = \sum_{\alpha\le\nu\le\beta}\frac{e(-\phi(\nu)-1/8)}{|f''(x_\nu|^{1/2}} + O(\log(FN^{-1}+2) + F^{-1/2}N). \tag{3.2}$$

Lemma 3. *(van der Corput's bound, Theorem 2.2 of [1]) Suppose that f is a real valued function with two continuous derivatives on $[a,b]$ where $a<b$ are integers. Suppose also that there is some $\lambda>0$ and some $\alpha\ge 1$ such that*

$$\lambda \le |f''(x)| \le \alpha\lambda$$

on $[a,b]$. Then

$$\sum_{a\le n\le b} e(f(n)) \ll \alpha(b-a)\lambda^{1/2} + \lambda^{-1/2}. \tag{3.3}$$

4. Estimation of the double exponential sum

We start with the integral positive definite quadratic form $Q^*(x,y)$ (see equation (3.4) of [2]) :

$$Q^*(x,y) = a^*x^2 + b^*xy + c^*y^2 \tag{4.1}$$

and let

$$d := (b^*)^2 - 4a^*c^* < 0.$$

Note that $a^* > 0, c^* > 0$. As remarked in [2], the form $Q^*(x,y)$ depends only on Q and k, and $|d| \leq \Delta$.

Our goal is to bound non-trivially the exponential sum (equation (4.4) of [2]):

$$\sum_x \sum_{y \in \mathcal{I}(x)} e\left(Q^*(x,y)\cdot r + \frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right)\right), \tag{4.2}$$

where

$$r := \frac{\overline{h\Delta_0}}{k} - \frac{1}{2hk\Delta_0}$$

and

$$\mathcal{I}(x) := \{y \in \mathbb{R} \ : \ N \leq Q^*(x,y) \leq N'\},$$

with the convention that the sums are restricted to residue classes mod $|d|$. Here, $1 \leq N \leq N' \leq 2N, N \ll K, \Delta_0|\Delta, h/k$ is a rational number satisfying $(h\Delta_0, k) = 1$ and

$$\frac{K}{T} \ll| \frac{1}{\sqrt{\Delta}} - \frac{h}{k} |\leq \frac{\pi K}{T\Delta},$$

$$k, h \asymp \sqrt{T/K}, \quad t \asymp T.$$

The function $\phi(x)$ is defined as

$$\phi(x) = \operatorname{arsinh}(x^{1/2}) + (x + x^2)^{1/2}.$$

In [2], the summation over $y-$ range was evaluated using a classical estimate for exponential sums (Lemma 4.1 of [2]). In the present paper, we explicitly carry out Weyl shift, apply B process and then exploit the cancellations in the sum over x invoking van der Corput's bound.

For the ease of exposition, we restrict ourselves to the case $|d| = 1$, i.e., x and y vary over all integers. The general case can be treated in a similar way.

It is easy to see that $\mathcal{I}(x)$ is empty unless $x \in \mathcal{J}$, where

$$\mathcal{J} := \left[-\frac{2\sqrt{c^*N'}}{\sqrt{|d|}}, \frac{2\sqrt{c^*N'}}{\sqrt{|d|}}\right]$$

and that

$$\mathcal{I}(x) = I(x) \cup I'(x),$$

where

$$I(x) := \left[\sqrt{\max\left\{0, \frac{N}{c^*} - \frac{|d|}{(2c^*)^2}\cdot x^2\right\}}, \sqrt{\frac{N'}{c^*} - \frac{|d|}{(2c^*)^2}\cdot x^2}\right]$$

and

$$I'(x) := \left[-\sqrt{\frac{N'}{c^*} - \frac{|d|}{(2c^*)^2}\cdot x^2}, -\sqrt{\max\left\{0, \frac{N}{c^*} - \frac{|d|}{(2c^*)^2}\cdot x^2\right\}}\right].$$

Set

$$J = \{x \in \mathcal{J} \ : \ x \geq 0\} = \left[0, \frac{2\sqrt{c^*N'}}{\sqrt{|d|}}\right].$$

In the following, we estimate the partial sum

$$\sum_{x\in J}\sum_{y\in I(x)} e\left(Q^*(x,y)\cdot r+\frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right)\right),$$

where x is positive and y runs over the interval $I(x)$. The remaining three partial sums with a) $x\geq 0$ and $y\in I'(x)$, b) $x<0$ and $y\in I(x)$, c) $x<0$ and $y\in I'(x)$ can be estimated in a similar way.

We start with applying Cauchy-Schwarz, getting

$$\begin{aligned}&\left|\sum_{x\in J}\sum_{y\in I(x)} e\left(Q^*(x,y)\cdot r+\frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right)\right)\right|^2\\ \ll&\sqrt{N}\cdot\sum_{x\in J}\left|\sum_{y\in I(x)} e\left(Q^*(x,y)\cdot r+\frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right)\right)\right|^2,\end{aligned}\tag{4.3}$$

where we use $|J|\ll\sqrt{N}$. Applying Lemma 1 with $\lambda=2$, and using $|J|\ll\sqrt{N}$ and $|I(x)|\ll\sqrt{N}$, we have

$$\begin{aligned}&\sum_{x\in J}\left|\sum_{y\in I(x)} e\left(Q^*(x,y)\cdot r+\frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right)\right)\right|^2\\ \ll&\sum_{x\in J}\frac{\sqrt{N}}{M}\cdot\sum_{0\leq|m|\leq M}\left(1-\frac{|2m|}{M}\right)\cdot\sum_{y\in I_m(x)} e\left(f_x(y+2m)-f_x(y)\right)\\ \ll&\frac{\sqrt{N}}{M}\cdot\sum_{1\leq m\leq M}\left|\sum_{x\in J}\sum_{y\in I(x)} e\left(f_x(y+m)-f_x(y-m)\right)\right|+NM+\frac{N^{3/2}}{M},\end{aligned}\tag{4.4}$$

where $M\leq N$ is any natural number,

$$I_m(x):=\{y\in I(x)\ :\ y+2m\in I(x)\}$$

and

$$f_x(y):=Q^*(x,y)\cdot r+\frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right).$$

We have

$$Q^*(x,y+m)-Q^*(x,y-m)=2m\left(b^*x+2c^*y\right)$$

and, using Taylor series expansion,

$$\phi\left(\frac{\pi Q^*(x,y+m)}{2hk\Delta_0 t}\right)-\phi\left(\frac{\pi Q^*(x,y-m)}{2hk\Delta_0 t}\right)=2mg_x'(y)+O\left(m^3\left|g_x'''(y)\right|\right),$$

where

$$g_x(y):=\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right).$$

Using

$$\phi'(u)=\frac{1}{\sqrt{u}}+O(\sqrt{u}),\quad \phi''(u)=O\left(\frac{1}{u^{3/2}}\right)\quad\text{and}\quad \phi'''(u)=O\left(\frac{1}{u^{5/2}}\right)$$

for $|u|\leq 1$, we calculate that

$$g_x'(y)=\frac{\pi\left(b^*x+2c^*y\right)}{2hk\Delta_0 t}\cdot\left(\frac{1}{\sqrt{\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}}}+O\left(\sqrt{\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}}\right)\right)=\frac{\sqrt{\pi}\left(b^*x+2c^*y\right)}{\sqrt{2hk\Delta_0 tQ^*(x,y)}}+O\left(\frac{NK^{3/2}}{T^3}\right)$$

and

$$g_x'''(y)=O\left(\frac{K^{1/2}}{NT}\right).$$

It follows that

$$\sum_{y\in I(x)} e\left(f_x(y+m)-f_x(y-m)\right) = \sum_{y\in I(x)} e\left(2mF_x(y)\right) + O\left(\frac{mN^{3/2}K^{3/2}}{T^2}+\frac{m^3K^{1/2}}{N^{1/2}}\right), \quad (4.5)$$

where

$$F_x(y) := (b^*x+2c^*y)\cdot\left(r+\frac{\sqrt{t}}{\sqrt{2\pi hk\Delta_0 Q^*(x,y)}}\right).$$

We calculate that

$$\begin{aligned} F_x'(y) =& 2c^*r+\frac{\sqrt{t}|d|x^2}{2\sqrt{2\pi hk\Delta_0}Q^*(x,y)^{3/2}},\\ F_x''(y) =& -\frac{3\sqrt{t}|d|x^2\left(b^*x+2c^*y\right)}{4\sqrt{2\pi hk\Delta_0}Q^*(x,y)^{5/2}} \asymp \frac{K^{1/2}}{N},\\ F_x'''(y) =& O\left(\frac{K^{1/2}}{N^{3/2}}\right),\\ F_x''''(y) =& O\left(\frac{K^{1/2}}{N^2}\right). \end{aligned} \quad (4.6)$$

By our assumptions on x and y, we have $N\le Q^*(x,y)\le N'$. For fixed x, we also have

$$Q^*(x,y)\ge \frac{|d|}{4c^*}\cdot x^2.$$

Hence $a(x)\le F_x'(y)\le b(x)$, where

$$a(x) := 2c^*r+\frac{\sqrt{t}|d|x^2}{2\sqrt{2\pi hk\Delta_0}N'^{3/2}}$$

and

$$\begin{aligned} b(x) :=& 2c^*r+\frac{\sqrt{t}|d|x^2}{2\sqrt{2\pi hk\Delta_0}\max\{N,|d|x^2/(4c^*)\}^{3/2}}\\ =& 2c^*r+\min\left\{\frac{\sqrt{t}|d|x^2}{2\sqrt{2\pi hk\Delta_0}N^{3/2}},\frac{\left(4c^*\right)^{3/2}\sqrt{t}}{2\sqrt{2\pi|d|hk\Delta_0}x}\right\}. \end{aligned}$$

Now Lemma 2 yields

$$\begin{aligned} &\sum_{y\in I(x)} e\left(2mF_x(y)\right)\\ =&\sum_{2ma(x)\le n\le 2mb(x)} \frac{e\left(2mF_x\left(y_{x,m,n}\right)-ny_{x,m,n}-1/8\right)}{\sqrt{2m|F_x''(y_{x,m,n})|}}+O\left(\log T+\frac{N^{1/2}}{m^{1/2}K^{1/4}}\right), \end{aligned} \quad (4.7)$$

where $y_{x,m,n}\in I(x)$ is the solution of $2mF_x'(y_{x,m,n})=n$.

We compute that

$$F_x\left(y_{x,m,n}\right) = \sqrt{-|d|x^2+4c^*\left(\frac{m\sqrt{t}|d|x^2}{\sqrt{2\pi hk\Delta_0}\left(n-4mc^*r\right)}\right)^{2/3}}\cdot\left(r+\left(\frac{\left(n-4mc^*r\right)t}{2\pi hk\Delta_0|d|mx^2}\right)^{1/3}\right)$$

and

$$ny_{x,m,n} = \frac{n}{2c^*}\cdot\left(-b^*x+\sqrt{-|d|x^2+4c^*\left(\frac{m\sqrt{t}|d|x^2}{\sqrt{2\pi hk\Delta_0}\left(n-4mc^*r\right)}\right)^{2/3}}\right).$$

Putting together gives

$$\begin{aligned}
&G_{m,n}(x) := 2mF_x\left(y_{x,m,n}\right) - ny_{x,m,n}\\
=&\frac{b^*n}{2c^*}\cdot x - \frac{1}{2c^*}\cdot\sqrt{-|d|x^2 + 4c^*\left(\frac{m\sqrt{t}|d|x^2}{\sqrt{2\pi hk\Delta_0}\left(n-4mc^*r\right)}\right)^{2/3}}\times\\
&\left(\left(n-4c^*mr\right) - 4c^*m\left(\frac{\left(n-4mc^*r\right)t}{2\pi hk\Delta_0|d|mx^2}\right)^{1/3}\right) \qquad (4.8)\\
=&\frac{b^*n}{2c^*}\cdot x + \frac{\left(n-4c^*mr\right)}{2c^*|d|x^2}\cdot\left(-|d|x^2 + 4c^*\left(\frac{m\sqrt{t}|d|x^2}{\sqrt{2\pi hk\Delta_0}\left(n-4mc^*r\right)}\right)^{2/3}\right)^{3/2}\\
=&\frac{b^*n}{2c^*}\cdot x + \frac{1}{2c^*}\cdot\left(-(n-4c^*mr)^{2/3}|d|^{1/3}x^{2/3} + 4c^*\cdot\frac{m^{2/3}t^{1/3}}{(2\pi hk\Delta_0)^{1/3}}\right)^{3/2}.
\end{aligned}$$

Combining (4.4), (4.5), (4.6), (4.7) and (4.8), we get

$$\begin{aligned}
&\left|\sum_{x\in J}\sum_{y\in I(x)} e\left(Q^*(x,y)\cdot r + \frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right)\right)\right|^2\\
\ll&\frac{N}{M}\cdot\sum_{1\le m\le M}\left|\sum_{x\in J}\sum_{ma(x)\le n\le mb(x)}\frac{e\left(G_{m,n}(x)\right)}{\sqrt{|F_x''(y_{x,m,n})|}}\right| + \qquad (4.9)\\
&O\left(MN^{3/2} + \frac{N^2}{M} + \frac{MN^{5/2}K^{3/2}}{T^2} + M^3N^{1/2}K^{1/2} + \frac{N^2}{M^{1/2}K^{1/4}}\right).
\end{aligned}$$

Let's first work out what the trivial estimate for the double exponential sum above gives. We have

$$|J| \ll N^{1/2} \quad \text{and} \quad b(x) - a(x) = O\left(K^{1/2}N^{-1/2}\right).$$

Together with (4.6), this implies

$$\frac{N}{M}\cdot\sum_{1\le m\le M}\left|\sum_{x\in J}\sum_{2ma(x)\le n\le 2mb(x)}\frac{e\left(G_{m,n}(x)\right)}{\sqrt{2m|F_x''(y_{x,m,n})|}}\right| = O\left(M^{1/2}N^{3/2}K^{1/4}\right). \qquad (4.10)$$

So choosing $M := \left[N^{1/6}\right]$, using $N \ll K$, and taking the square root, we deduce that

$$\begin{aligned}
&K^{1/4}N^{-1/4}T^{-1/2}\left|\sum_{x\in J}\sum_{y\in I(x)} e\left(Q^*(x,y)\cdot r + \frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right)\right)\right| \qquad (4.11)\\
=&O\left(\frac{K^{11/12}}{T^{1/2}} + \frac{K^{25/12}}{T^{3/2}}\right).
\end{aligned}$$

So if

$$K := T^{6/11-\varepsilon},$$

the above is $O\left(T^{-\varepsilon}\right)$, as desired.

Now we estimate the said double exponential sum non-trivially. First, we re-arrange summations, getting

$$\sum_{x\in J}\sum_{2ma(x)\le n\le 2mb(x)}\frac{e\left(G_{m,n}(x)\right)}{\sqrt{2m|F_x''(y_{x,m,n})|}} = \sum_{n\in J_m}\sum_{A_m(n)\le x\le B_m(n)}\frac{e\left(G_{m,n}(x)\right)}{\sqrt{2m|F_x''(y_{x,m,n})|}}, \qquad (4.12)$$

where

$$J_m = \left[4c^*mr, 4c^*mr + \frac{2c^*m\sqrt{t}N'}{\sqrt{2\pi hk\Delta_0}N^{3/2}}\right],$$

$$A_m(n) := (n-4mc^*r)^{1/2} \cdot \frac{(2\pi hk\Delta_0)^{1/4}N^{3/4}}{(m|d|)^{1/2}t^{1/4}}$$

and

$$B_m(n) := \min\left\{ \frac{2(c^*N')^{1/2}}{|d|^{1/2}}, (n-4mc^*r)^{1/2} \cdot \frac{(2\pi hk\Delta_0)^{1/4}N'^{3/4}}{(m|d|)^{1/2}t^{1/4}}, \frac{(4c^*)^{3/2}\,mt^{1/2}}{(2\pi|d|hk\Delta_0)^{1/2}\,(n-4mc^*r)} \right\}.$$

We note that

$$|J_m| \ll \frac{mK^{1/2}}{N^{1/2}} \quad \text{and} \quad B_m(n) - A_m(n) \ll N^{1/2} \tag{4.13}$$

and compute that

$$\begin{aligned} G''_{m,n}(x) = &- \frac{(n-4c^*mr)^{2/3}|d|^{1/3}}{6c^*x^{4/3}} \cdot \left((n-4c^*mr)^{2/3}|d|^{1/3}x^{2/3} + 4c^* \cdot \frac{m^{2/3}t^{1/3}}{(2\pi hk\Delta_0)^{1/3}} \right) \times \\ &\left(-(n-4c^*mr)^{2/3}|d|^{1/3}x^{2/3} + 4c^* \cdot \frac{m^{2/3}t^{1/3}}{(2\pi hk\Delta_0)^{1/3}} \right)^{-1/2}. \end{aligned} \tag{4.14}$$

We have

$$\frac{(n-4c^*mr)^{2/3}|d|^{1/3}}{6c^*x^{4/3}} \asymp \frac{m^{2/3}t^{1/3}}{(hk\Delta_0)^{1/3}N} \asymp \frac{m^{2/3}K^{1/3}}{N}$$

and

$$(n-4c^*mr)^{2/3}|d|^{1/3}x^{2/3} + 4c^* \cdot \frac{m^{2/3}t^{1/3}}{(2\pi hk\Delta_0)^{1/3}} \asymp \frac{m^{2/3}t^{1/3}}{(hk\Delta_0)^{1/3}} \asymp m^{2/3}K^{1/3}$$

and hence

$$G''_{m,n}(x) \asymp \frac{m^{4/3}K^{2/3}}{N} \cdot \left(-(n-4c^*mr)^{2/3}|d|^{1/3}x^{2/3} + 4c^* \cdot \frac{m^{2/3}t^{1/3}}{(2\pi hk\Delta_0)^{1/3}} \right)^{-1/2} \tag{4.15}$$

for n and x in the relevant summation intervals.

We first assume that $n \in J'_m \subseteq J_m$, where

$$J'_m := \left[4c^*mr, 4c^*mr + \frac{c^*m\sqrt{t}}{\sqrt{2\pi hk\Delta_0 N'}} \right),$$

in which case we compute that

$$-(n-4c^*mr)^{2/3}|d|^{1/3}x^{2/3} + 4c^* \cdot \frac{m^{2/3}t^{1/3}}{(2\pi hk\Delta_0)^{1/3}} \asymp \frac{m^{2/3}t^{1/3}}{(hk\Delta_0)^{1/3}} \asymp m^{2/3}K^{1/3}$$

and hence, using (4.15),

$$G''_{m,n}(x) \asymp \frac{mK^{1/2}}{N} \quad \text{if } A_m(n) \le x \le B_m(n). \tag{4.16}$$

Now we apply partial summation to remove the factor $1/\sqrt{F''_x(y_{x,m,n})}$ and Lemma 3 to get

$$\begin{aligned} &\sum_{n\in J'_m} \sum_{A_m(n)\le x\le B_m(n)} \frac{e\left(G_{m,n}(x)\right)}{\sqrt{2m|F''_x(y_{x,m,n})|}} \\ \ll &\frac{N^{1/2}}{m^{1/2}K^{1/4}} \cdot \left(mK^{1/2} \cdot \left(\frac{mK^{1/2}}{N}\right)^{1/2} + \frac{mK^{1/2}}{N^{1/2}} \cdot \left(\frac{mK^{1/2}}{N}\right)^{-1/2} \right) \ll mK^{1/2}, \end{aligned} \tag{4.17}$$

where we have used (4.13) and (4.16).

Next, we assume that $n \in J''_m \subseteq J_m$, where

$$J''_m := \left[4c^*mr + \frac{c^*m\sqrt{t}}{\sqrt{2\pi hk\Delta_0 N'}}, 4c^*mr + \frac{2c^*m\sqrt{t}N'}{\sqrt{2\pi hk\Delta_0}N^{3/2}} \right] \tag{4.18}$$

and note that in this case

$$x \asymp N^{1/2} \quad \text{if } A_m(n) \le x \le B_m(n). \tag{4.19}$$

We further note that

$$J_m = J'_m \cup J''_m. \tag{4.20}$$

We write

$$I_m(n) := \left(\frac{(4c^*)^{3/2} mt^{1/2}}{(2\pi|d|hk\Delta_0)^{1/2}(n-4mc^*r)} - 1, \frac{(4c^*)^{3/2} mt^{1/2}}{(2\pi|d|hk\Delta_0)^{1/2}(n-4mc^*r)} \right]$$

and

$$I_m(n,\delta) := \left(\frac{(4c^*)^{3/2} mt^{1/2}}{(2\pi|d|hk\Delta_0)^{1/2}(n-4mc^*r)} - 2\delta, \frac{(4c^*)^{3/2} mt^{1/2}}{(2\pi|d|hk\Delta_0)^{1/2}(n-4mc^*r)} - \delta \right]$$

and set

$$S_m(n) := I_m(n) \cap [A_m(n), B_m(n)] \quad \text{and} \quad S_m(n,\delta) := I_m(n,\delta) \cap [A_m(n), B_m(n)].$$

We observe that the interval $[A_m(n), B_m(n)]$ can be split into $O(\log T)$ intervals of the form $S_m(n)$ or $S_m(n,\delta)$, where

$$1 \ll \delta \ll \sqrt{N}. \tag{4.21}$$

Estimating trivially gives

$$\sum_{n\in J_m''} \sum_{x\in S_m(n)} \frac{e(G_{m,n}(x))}{\sqrt{2m|F_x''(y_{x,m,n})|}} \ll m^{1/2}K^{1/4}, \tag{4.22}$$

where we have used (4.6) and (4.13). If $x \in S_m(n,\delta)$, then we compute using (4.18), (4.19) and the mean value theorem that

$$-(n-4c^*mr)^{2/3}|d|^{1/3}x^{2/3} + 4c^* \cdot \frac{m^{2/3}t^{1/3}}{(2\pi hk\Delta_0)^{1/3}} \asymp \delta \cdot \left(\frac{m\sqrt{t}}{\sqrt{hk\Delta_0}N^{1/2}} \right)^{2/3} x^{-1/3} \asymp \frac{\delta m^{2/3}K^{1/3}}{N^{1/2}}.$$

Hence, using (4.15), it follows that

$$G_{m,n}''(x) \asymp \frac{mK^{1/2}}{\delta^{1/2}N^{3/4}}. \tag{4.23}$$

Again using partial summation to remove the factor $1/\sqrt{F_x''(y_{x,m,n})}$ and Lemma 3, we deduce that

$$\begin{aligned} &\sum_{n\in J_m''} \sum_{x\in S_m(n,\delta)} \frac{e(G_{m,n}(x))}{\sqrt{2m|F_x''(y_{x,m,n})|}} \\ &\ll \frac{N^{1/2}}{m^{1/2}K^{1/4}} \cdot \left(\frac{\delta mK^{1/2}}{N^{1/2}} \cdot \left(\frac{mK^{1/2}}{\delta^{1/2}N^{3/4}} \right)^{1/2} + \frac{mK^{1/2}}{N^{1/2}} \cdot \left(\frac{mK^{1/2}}{\delta^{1/2}N^{3/4}} \right)^{-1/2} \right) \ll mK^{1/2}, \end{aligned} \tag{4.24}$$

where we have employed $|S_m(n,\delta)| \le \delta$, (4.13), (4.21) and (4.23). From (4.22) and (4.24), it follows that

$$\sum_{n\in J_m''} \sum_{A_m(n)\le x\le B_m(n)} \frac{e(G_{m,n}(x))}{\sqrt{2m|F_x''(y_{x,m,n})|}} \ll mK^{1/2}\log T. \tag{4.25}$$

Now combining (4.17), (4.20) and (4.25), we get

$$\sum_{n\in J_m} \sum_{A_m(n)\le x\le B_m(n)} \frac{e(G_{m,n}(x))}{\sqrt{2m|F_x''(y_{x,m,n})|}} \ll mK^{1/2}\log T. \tag{4.26}$$

Combining (4.9), (4.12) and (4.26), we deduce that

$$\begin{aligned} &\left| \sum_{x\in J} \sum_{y\in I(x)} e\left(Q^*(x,y)\cdot r + \frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t} \right) \right) \right|^2 \\ &= O\left(MN^{3/2} + \frac{N^2}{M} + \frac{MN^{5/2}K^{3/2}}{T^2} + M^3N^{1/2}K^{1/2} + \frac{N^2}{M^{1/2}K^{1/4}} + MNK^{1/2}\log T \right). \end{aligned} \tag{4.27}$$

Choosing $M := \left[N^{1/4}\right]$, using $N \ll K$, and taking the square root, it follows that

$$\begin{aligned} &K^{1/4}N^{-1/4}T^{-1/2}\left|\sum_{x\in J}\sum_{y\in I(x)} e\left(Q^*(x,y)\cdot r + \frac{t}{\pi}\cdot\phi\left(\frac{\pi Q^*(x,y)}{2hk\Delta_0 t}\right)\right)\right| \\ =&O\left(\frac{K^{7/8}}{T^{1/2}}\log T + \frac{K^{17/8}}{T^{3/2}}\right). \end{aligned}$$

This is $O(T^{-\varepsilon})$, provided that $K := T^{4/7-\varepsilon}$, which completes the proof.

Acknowledgement. The third author would like to thank the Institute of Mathematical Sciences, Chennai for supporting her visits which enabled her to work on this project.

References

[1] S. W. Graham and G. Kolesnik, Van der Corput's Method of Exponential sums, *Cambridge University Press*, New York, 1991.
[2] M. Jutila and K. Srinivas, Gaps between the zeros of Epstein's zeta-functions on the critical line, *Bull. London Math. Soc.* **37** (2005) 45-53.
[3] H. S. A. Potter and E. C. Titchmarsh, The zeros of Epstein's zeta-functions, *Proc. London Math. Soc.* **39**(2) (1935) 372–384.
[4] A. Sankaranarayanan, Zeros of quadratic zeta-functions on the critical line, *Acta Arith.* **69** (1995) 21–37.

IISER-Thiruvananthapuram, CET Campus, Trivandrum-695016, Kerala, India

Institute of Mathematical Sciences, CIT Campus, Taramani, Chennai 600 113, India

SRTM University, Nanded, Maharashtra 431606, India
E-mail address, Stephan Baier: sbaier@math.tifr.res.in
E-mail address, Kotyada Srinivas: srini@imsc.res.in
E-mail address, Usha Sangale: ushas073@gmail.com